arXiv:1402.0947v4 [stat.OT] 22 Apr 2014

On Rényi entropy convergence of the max domain of attraction

Ali Saeb[1]
Theoretical Statistics and Mathematics Unit,
Indian Statistical Institute, Delhi Center,
7 S.J.S Sansanwal Marg, New Delhi 110016, India

Abstract: In this paper, we prove that the Rényi entropy of linearly normalized partial maxima of independent and identically distributed random variables is convergent to the corresponding limit Rényi entropy when the linearly normalized partial maxima converges to some nondegenerate random variable.

Keywords: Rényi entropy, Max stable laws, Max domain of attraction.

MSC 2010 classification: 60F10

[1]Corresponding author: ali.saeb@gmail.com

1. Introduction

The limit laws of linearly normalized partial maxima $M_n = \max(X_1, \cdots, X_n)$ of independent and identically distributed (iid) random variables (rvs) $X_1, X_2, \ldots$, with common distribution function (df) F, namely,

$$\lim_{n\to\infty} \Pr(M_n \le a_n x + b_n) = \lim_{n\to\infty} F^n(a_n x + b_n) = G(x), \quad x \in \mathcal{C}(G), \tag{1.1}$$

where, $a_n > 0$, $b_n \in \mathbb{R}$, are norming constants, G is a non-degenerate distribution function, $\mathcal{C}(G)$ is the set of all continuity points of G, are called max stable laws. If, for some non-degenerate distribution function G, a distribution function F satisfies (1.1) for some norming constants $a_n > 0$, $b_n \in \mathbb{R}$, then we say that F belongs to the max domain of attraction of G under linear normalization and denote it by $F \in \mathcal{D}(G)$. Limit distribution functions G satisfying (1.1) are the well known extreme value types of distributions, or max stable laws, namely,

$$\text{the Fréchet law:} \quad \Phi_\alpha(x) = \begin{cases} 0, & x < 0, \\ \exp(-x^{-\alpha}), & 0 \le x; \end{cases}$$

$$\text{the Weibull law:} \quad \Psi_\alpha(x) = \begin{cases} \exp(-|x|^\alpha), & x < 0, \\ 1, & 0 \le x; \end{cases}$$

$$\text{and the Gumbel law:} \quad \Lambda(x) = \exp(-\exp(-x)); \quad x \in \mathbb{R};$$

$\alpha > 0$ being a parameter, with respective density functions,

$$\text{the Fréchet density:} \quad \phi_\alpha(x) = \begin{cases} 0, & x \le 0, \\ \alpha x^{-(\alpha+1)} e^{-x^{-\alpha}}, & 0 < x; \end{cases}$$

$$\text{the Weibull density:} \quad \psi_\alpha(x) = \begin{cases} \alpha |x|^{\alpha-1} e^{-|x|^\alpha}, & x < 0, \\ 0, & 0 \le x; \end{cases}$$

$$\text{and the Gumbel density:} \quad \lambda(x) = e^{-x} e^{-e^{-x}}, \quad x \in \mathbb{R}.$$

Note that (1.1) is equivalent to

$$\lim_{n\to\infty} n(1 - F(a_n x + b_n)) = -\log G(x), \ x \in \{y : G(y) > 0\}.$$

We shall denote the left extremity of distribution function F by $l(F) = \inf\{x : F(x) > 0\} \ge -\infty$ and the right extremity of F by $r(F) = \sup\{x : F(x) < 1\} \le \infty$. Criteria for $F \in \mathcal{D}(G)$ are well known (see, for example, Galambos, 1987; Resnick, 1987; Embrechts et al., 1997).

The Shannon entropy of a continuous rv X with density function $f(x)$ is defined as

$$H(X) = -\int_A f(x) \log f(x) dx, \quad \text{where} \ \ A = \{x \in \mathbb{R} : f(x) > 0\}.$$

Rényi entropy is a generalization of Shannon entropy (Rényi, 1961). It is one of a family of functional for quantifying the diversity, uncertainty or randomness of a system. The Rényi entropy of order β is defined as

$$H_\beta(X) = \frac{1}{1-\beta} \log \left(\int_A (f(x))^\beta \, dx \right), \tag{1.2}$$

where, $0 < \beta < \infty$, $\beta \ne 1$. By L'Hopital's rule, the Rényi entropy tends to Shannon entropy, as $\beta \to 1$. The Rényi entropy are important in ecology and statistics as indices of diversity. Rényi entropy appear also in several important contexts such as those of information theory, statistical estimation.

The idea of tracking the central limit theorem using Shannon entropy goes back to Linnik (1959) and Shimizu (1975), who used it to give a particular proof of the central limit theorem. Brown (1982), Barron (1986) and Takano (1987) discuss the central limit theorem with convergence in the sense of Shannon entropy and relative entropy. Artstein et al.(2004) and Johnson and Barron (2004) obtained the rate of convergence under some conditions on

the density. Johnson (2006) is a good reference to the application of information theory to limit theorems, especially the central limit theorem. Cui and Ding (2010) show that the convergence of the Rényi entropy of the normalized sums of iid rvs and obtain the corresponding rates of convergence. Saeb (2014) study the rate of convergence of Rényi entropy for the max domain of attraction.

In this article, our main interest is to investigate conditions under which the Rényi entropy of the normalized partial maxima of iid rvs converges to the corresponding limit Rényi entropy. In the other hand, our problem of interest to see if normalized partial maxima converges to a nondegenerate rv, does the Rényi entropy of the normalized partial maxima converges to the Rényi entropy of the limit rv? In the next section we give our main results, followed by a section on Proofs. The Rényi entropies of the extreme value distributions are given in the appendix A with proof and appendix B containing results used in this article.

2. Main Result

Theorem 2.1. Suppose $F \in D(G)$ is absolutely continuous with pdf f which is eventually positive and decreasing in left neighbourhood of $r(F)$. If $\int_{-\infty}^{r(F)} (f(x))^{\beta} dx < \infty$ for $\beta > 1$, and

(a) $G = \Phi_\alpha$ and $r(F) = \infty$, then $\lim_{n\to\infty} H_\beta(g_n) = H_\beta(\phi_\alpha)$;

(b) $G = \Psi_\alpha$ and $r(F) < \infty$, then $\lim_{n\to\infty} H_\beta(g_n) = H_\beta(\psi_\alpha)$;

(c) $G = \Lambda$ and $r(F) \leq \infty$, then $\lim_{n\to\infty} H_\beta(g_n) = H_\beta(\lambda)$.

Remark 2.1. Lemma A.2 show that, the Rényi entropy do not depend on the location and scale parameters.

The proof of the above theorem is different from the proof for the Rényi entropy of the normalized sums of iid rvs. In our proofs, the properties of normalized partial maxima such as, von Mises condition and density convergence, plays an important role.

3. Proofs

Lemma 3.1. If $\bar{F} \in RV_{-\alpha}$ then $a(\cdot) \in RV_{1/\alpha}$.

Proof. We know that $1 - F$ is regularly varying so that $n\bar{F}(a_n x) \to x^{-\alpha}$ as $n \to \infty$. Set $U = 1/\bar{F}$ and $V = U^{\leftarrow}$. Therefore

$$U(a_n x)/n \to x^{\alpha}, \text{ for } x > 0,$$

and inverting we have

$$V(ny)/a_n \to y^{-1/\alpha}.$$

Since, $a_n \simeq (\frac{1}{1-F})^{\leftarrow}(n) = V(n)$ and a switch to a continuous variable we have

$$\frac{V(ty) - V(t)}{a(t)} \to y^{1/\alpha} - 1.$$

Now we have,

$$\begin{aligned} \lim_{t\to\infty} \frac{a(tx)}{a(t)} &= \lim_{t\to\infty} \frac{a(tx)(V(tx) - V(t))}{-a(t)(V(tx\,x^{-1} - V(tx)))}, \\ &= (x^{1/\alpha} - 1)/(1 - x^{-1/\alpha}), \\ &= x^{1/\alpha}. \end{aligned}$$

□

Proof of Theorem 2.1-(a). Suppose $F \in \mathcal{D}(\Phi_\alpha)$, and $1-F$ is regularly varying so that

$$\begin{aligned} \lim_{n\to\infty} \frac{\overline{F}(a_n x)}{\overline{F}(a_n)} &= x^{-\alpha}, \ \ x>0; \ \text{ and} \\ \lim_{n\to\infty} F^n(a_n x) &= \Phi_\alpha(x), \ x \in \mathbb{R}, \end{aligned}$$

with $a_n = F^{\leftarrow}(1-\frac{1}{n}) = \inf\{x : F(x) > 1-\frac{1}{n}\}$, $n \geq 1$ and $b_n = 0$. From Theorem B.4, F satisfies the von Mises condition:

$$\lim_{t\to\infty} \frac{tf(t)}{1-F(t)} = \alpha. \tag{3.1}$$

Now, by Theorem B.6, implies the following density convergence on compact sets:

$$\lim_{n\to\infty} g_n(x) = \phi_\alpha(x), \ \ x \in K \subset (0,\infty) \tag{3.2}$$

where K is a compact set, and $g_n(x) = na_n f(a_n x)F^{n-1}(a_n x)$.

From definition of Rényi entropy, we write,

$$H_\beta(g_n) = \frac{1}{1-\beta}\log\left[I_A(n,v) + I_B(n,v) + I_C(n,v)\right]. \tag{3.3}$$

where, $I_A(n,v) = \int_v^\infty (g_n(x))^\beta\, dx$, and $I_B(n,v) = \int_{-\infty}^{v^{-1}} (g_n(x))^\beta\, dx$, and $I_C(n,v) = \int_{v^{-1}}^{v} (g_n(x))^\beta\, dx$. It is enough to show that

$$\lim_{v\to\infty}\lim_{n\to\infty} (I_A(n,v) + I_B(n,v)) = 0.$$

We set,

$$\begin{aligned} 0 < I_A(n,v) &= \int_v^\infty (g_n(x))^{\beta-1}\, dF^n(a_n x), \\ &= L(n,\beta)\int_v^\infty \left(\frac{f(a_n x)}{f(a_n)} F^{(n-1)}(a_n x)\right)^{\beta-1} g_n(x)dx, \\ &< L(n,\beta)\left(\frac{f(a_n v)}{f(a_n)}\right)^{\beta-1}(1-F^n(a_n v)), \end{aligned}$$

where, $L(n,\beta) = \left(\dfrac{a_n f(a_n) n\bar{F}(a_n)}{\bar{F}(a_n)}\right)^{\beta-1}$, and, $n\bar{F}(a_n) = 1$ and using von Mises conditions in (3.1) f is decreasing function, $L(n,\beta) \to \alpha^{\beta-1}$, as $n \to \infty$, and $\frac{f(a_n x)}{f(a_n)} < 1$ for $x \geq 1$. Hence, for $\beta > 1$,

$$\lim_{v\to\infty}\lim_{n\to\infty} I_A(n,v) = 0. \tag{3.4}$$

Now, we choose ξ_n by $-\log F(\xi_n) \simeq n^{-1/2}$, and $t_n = \frac{\xi_n}{a_n}$. If $\frac{\xi_n}{a_n} \to c > 0$ then $n^{1/2} \simeq -n(\log F(t_n\, a_n)) \to c^{-\alpha}$ and this is contradict the fact that $n^{1/2} \to \infty$. Therefore, $t_n \to 0$ as $\xi_n \to \infty$ for large n.
We have,

$$I_B(n,u) = \int_{-\infty}^{t_n} (g_n(x))^\beta dx + \int_{t_n}^{v^{-1}} (g_n(x))^\beta dx = I_{B_1}(n) + I_{B_2}(n,v). \tag{3.5}$$

We set,

$$\begin{aligned} I_{B_1}(n) &= n^{\beta} a_n^{\beta-1} \int_{-\infty}^{\xi_n} (F^{n-1}(s) f(s))^{\beta}\, ds, \ \ (\text{where, } a_n x = s), \\ &\leq n^{\beta} a_n^{\beta-1} F^{\beta(n-1)}(\xi_n) \int_{-\infty}^{\infty} (f(s))^{\beta}\, ds, \\ &\simeq \frac{n^{\beta} a_n^{\beta-1}}{\exp\{(n-1)\beta n^{-1/2}\}} \int_{-\infty}^{\infty} (f(s))^{\beta}\, ds, \end{aligned}$$

Since $a_n \simeq \left(\frac{1}{1-F}\right)^{\leftarrow}(n)$ from Lemma 3.1, $a_n \in RV_{\frac{1}{\alpha}}$ and (B.2) for $n > N$ given $\epsilon > 0$ and $\rho(n) < \frac{1+\epsilon}{\alpha}$ then

$$\begin{aligned} a_n &= c(n) \exp\Big\{ \int_N^n \rho(t) t^{-1} dt \Big\}, \\ &< (n/N)^{\frac{1+\epsilon}{\alpha}} c, \end{aligned}$$

Therefore,

$$0 < I_{B_1}(n) \leq \frac{c\, n^{\beta} (n/N)^{\frac{1+\epsilon}{\alpha}(\beta-1)}}{\exp\{\beta(n^{1/2} - n^{-1/2}\}} \int_{-\infty}^{\infty} (f(s))^{\beta}\, ds.$$

If $\int_{-\infty}^{\infty} (f(x))^{\beta} dx < \infty$ then,

$$\lim_{n\to\infty} I_{B_1}(n) = 0. \tag{3.6}$$

From (B.3), for given $\epsilon_1 > 0$ we have $f(a_n x) \leq -(\alpha+\epsilon_1)\frac{\log F(a_n x)}{a_n x}$ ultimately and from Theorem B.2 for sufficiently large n given $\epsilon_2 > 0$ and $n\bar{F}(a_n x) < \frac{1}{1-\epsilon_2} x^{-(\alpha-\epsilon_2)}$ and given $\epsilon_3 > 0$ and for all $n > 1$ then $-\frac{n-1}{n} < -(1-\epsilon_3)$. Hence, for sufficiently large n such that $n > n_0$ we have

$$\begin{aligned} g_n(x) &= n a_n f(a_n x) F^{n-1}(a_n x), \\ &< -(\alpha+\epsilon_1) n \log F(a_n x) x^{-1} \exp\Big\{ \frac{n-1}{n} n \log F(a_n x) \Big\}, \\ &< \frac{\alpha+\epsilon_1}{1-\epsilon_2} x^{-1-\alpha+\epsilon_2} \exp\Big\{ -\frac{1-\epsilon_3}{1-\epsilon_2} x^{-(\alpha-\epsilon_2)} \Big\}, \\ &< c\alpha' x^{-\alpha'-1} \exp\Big\{ -c x^{-\alpha'} \Big\}. \end{aligned}$$

where, $\alpha' = \alpha - \epsilon_2$, and c is positive constant. We define

$$h(x) = c\alpha' x^{-\alpha'-1} \exp\Big\{ -c x^{-\alpha'} \Big\}. \tag{3.7}$$

Set, $I_{B_2}(n, v) = \int_{t_n}^{v^{-1}} (g_n(x))^{\beta}\, dx$. From (3.7) for large n, $g_n(x) < h(x)$ we have

$$0 < I_{B_2}(n, v) < \int_0^{v^{-1}} (h(x))^{\beta} dx,$$

Since $\int_0^{\infty} (h(x))^{\beta} < \infty$ so that,

$$\lim_{v\to\infty} \lim_{n\to\infty} I_{B_2}(n, v) = 0. \tag{3.8}$$

From, (3.5), (3.6) and (3.8)

$$\lim_{v\to\infty} \lim_{n\to\infty} I_B(n, v) = 0. \tag{3.9}$$

Next, $I_C(n,v) = \int_{v^{-1}}^{v} (g_n(x))^\beta \, dx$, and from (3.7) for large n, we have $g_n(x) < h(x)$, and $\int_{v^{-1}}^{v} h(x)dx < \infty$, and using (3.2) $\lim_{n\to\infty} g_n(x) = \phi_\alpha(x)$, locally uniformly convergence in $x \in [v^{-1}, v]$ by DCT,

$$\lim_{v\to\infty}\lim_{n\to\infty} \int_{v^{-1}}^{v} (g_n(x))^\beta \, dx \quad = \quad \int_0^\infty (\phi_\alpha(x))^\beta \, dx. \tag{3.10}$$

And from (3.3), (3.4), (3.9) and (3.10) imply,

$$\lim_{n\to\infty} H_\beta(g_n) \quad = \quad H_\beta(\phi_\alpha).$$

□

Let $Y_1, Y_2, \ldots$ are iid rvs with common distribution function F_Y and $r(F_Y) < \infty$ and $X_i = 1/(r(F_Y) - Y_i)$ with common distribution function F_X. In following lemma is easily show that the relationship between the domain attraction of Φ_α and Ψ_α.

Lemma 3.2. If $F_Y \in \mathcal{D}(\Psi_\alpha)$ then $F_X \in \mathcal{D}(\Phi_\alpha)$ with $a_n = \frac{1}{\delta_n}$ and $b_n = 0$.

Proof. From $F_Y \in \mathcal{D}(\Psi_\alpha)$ with $\delta_n > 0$ and $r(F_Y) < \infty$, we have, $\frac{\vee_{i=1}^n Y_i - r(F_Y)}{\delta_n} \xrightarrow{d} M$, where, M is rv with distribution function Ψ_α. Therefore,

$$\begin{aligned} -\left(\frac{\vee_{i=1}^n Y_i - r(F_Y)}{\delta_n}\right)^{-1} &= \left(\vee_{i=1}^n \frac{\delta_n}{r(F_Y) - Y_i}\right), \\ &= \vee_{i=1}^n \frac{X_i}{a_n} \xrightarrow{d} -\frac{1}{M}, \end{aligned} \tag{3.11}$$

and $-\frac{1}{M}$ is a rv with distribution function Φ_α and left of (3.11) is equivalent to $F_X \in \mathcal{D}(\Phi_\alpha)$ with $a_n = \frac{1}{\delta_n}$ and $b_n = 0$. □

Proof of Theorem 2.1-(b). Suppose $F \in \mathcal{D}(\Psi_\alpha)$, iff $r(F) < \infty$ and $\overline{F}(r(F) - x^{-1}) \in RV_{-\alpha}$, is regularly varying. In this case we may set $\tau_n = F^{-1}(1 - \frac{1}{n})$, and $\delta_n = (r(F) - \tau_n)$, and then

$$\lim_{n\to\infty} F^n(\delta_n x + r(F)) = \Psi_\alpha(x), \; x \in \mathbb{R},$$

From Theorem B.4, F satisfies the von Mises condition:

$$\lim_{x\uparrow r(F)} \frac{(r(F) - x)f(x)}{\overline{F}(x)} = \alpha.$$

Now, by Theorem B.6, implies the following density convergence on compact sets:

$$\lim_{n\to\infty} g_n(x) = \psi_\alpha(x), \; x \in K \subset (-\infty, 0),$$

where K is a compact set, and $g_n(x) = n\delta_n f(\delta_n x + r(F))F^{n-1}(\delta_n x + r(F))$. From definition of Rényi entropy, we write,

$$H_\beta(g_n) \quad = \quad \frac{1}{1-\beta} \log\left(\int_{-\infty}^{0} (g_n(x))^\beta \, dx\right).$$

Put $x = -1/y$, and using Lemma 3.2, we have,

$$H_\beta(g_n) \quad = \quad \frac{1}{1-\beta} \log\left(\int_0^\infty (\tilde{g}_n(x))^\beta \frac{dy}{y^2}\right),$$

where, $\tilde{g}_n(x) = n\delta_n f(r(F) - \delta_n/y)F^{n-1}(r(F) - \delta_n/y)$. From Lemma 3.2, $F_X(a_n y) = F(r(F) - \delta_n/y) \in \mathcal{D}(\Phi_\alpha)$, and with conditions Theorem 2.1-(a), if $\int_{-\infty}^{r(F)} (f(s))^\beta ds < \infty$ and f is decreasing function for $\beta > 1$ then

$$\lim_{n\to\infty} H_\beta(g_n) \quad = \quad H_\beta(\psi_\alpha).$$

□

Lemma 3.3. Suppose $F \in \mathcal{D}(\Lambda)$ with auxiliary function u and $\epsilon > 0$. There exists a large N such that for $x \geq 0$, and $n > N$

$$n(1 - F(a_n x + b_n)) \leq (1+\epsilon)^2 (1 - \epsilon x)^{\epsilon^{-1}},$$

and for $x < 0$,

$$n(1 - F(a_n x + b_n)) \leq (1+\epsilon)^2 (1 + \epsilon |x|)^{\epsilon^{-1}}.$$

Proof. From Theorem B.8 and for $x \geq 0$, and sufficient large n such that $|u'(a_n t + b_n)| \leq \epsilon$

$$\begin{aligned} \left| \frac{u(a_n x + b_n)}{u(b_n)} - 1 \right| &= \left| \int_{b_n}^{a_n x + b_n} \frac{u'(s)}{u(b_n)} ds \right|, \\ &\leq \int_0^x |u'(a_n t + b_n)| \, dt, \quad \text{where } a_n s + b_n = t, \\ &\leq \epsilon \, x. \end{aligned} \tag{3.12}$$

Consequently, $\frac{u(b_n)}{u(a_n x + b_n)} < \frac{1}{1 - \epsilon \, x}$. Since, for large n, and $\epsilon > 0$ we have $\frac{(1-\epsilon)}{F(b_n)} < n < \frac{(1+\epsilon)}{F(b_n)}$ then

$$\begin{aligned} n(1 - F(a_n x + b_n)) &\leq (1+\epsilon) \frac{1 - F(b_n + a_n x)}{1 - F(b_n)}, \\ &= (1+\epsilon) \frac{c(b_n + a_n x)}{c(b_n)} \exp\Big\{ - \int_{b_n}^{b_n + a_n x} \frac{dy}{u(y)} \Big\}, \\ &\leq (1+\epsilon)^2 \exp\Big\{ - \int_0^x \frac{u(t) ds}{u(b_n + a_n s)} \Big\}, \end{aligned}$$

where, $y = b_n + a_n s$ and $\lim_{t \to \infty} \frac{c(t + x u(t))}{c(t)} = 1$ and from (3.12), then

$$\begin{aligned} n(1 - F(a_n x + b_n)) &\leq (1+\epsilon)^2 \exp\Big\{ - \int_0^x \frac{dt}{1 - \epsilon t} \Big\}, \\ &= (1+\epsilon)^2 (1 - \epsilon x)^{\epsilon^{-1}}. \end{aligned}$$

For the second statement, for $x < 0$ and sufficient large n

$$\begin{aligned} \left| 1 - \frac{u(a_n x + b_n)}{u(b_n)} \right| &= \left| \int_{a_n x + b_n}^{b_n} \frac{u'(s)}{u(b_n)} ds \right|, \\ &\leq \int_x^0 |u'(a_n t + b_n)| \, dt, \quad \text{where } a_n s + b_n = t, \\ &\leq \epsilon \, |x| \, . \end{aligned} \tag{3.13}$$

Consequently, $\frac{u(b_n)}{u(a_n x + b_n)} < \frac{1}{1 + \epsilon \, |x|}$. Then,

$$\begin{aligned} n(1 - F(a_n x + b_n)) &\leq (1+\epsilon) \frac{c(b_n + a_n x)}{c(b_n)} \exp\Big\{ - \int_{b_n}^{b_n + a_n x} \frac{dy}{u(y)} \Big\}, \\ &\leq (1+\epsilon)^2 \exp\Big\{ \int_x^0 \frac{u(t) ds}{u(b_n + a_n s)} \Big\}, \end{aligned}$$

where, $y = b_n + a_n s$ and $\lim_{t \to \infty} \frac{c(t + x u(t))}{c(t)} = 1$ and from (3.13), then

$$\begin{aligned} n(1 - F(a_n x + b_n)) &\leq (1+\epsilon)^2 \exp\Big\{ \int_x^0 \frac{dt}{1 + \epsilon \, |t|} \Big\}, \\ &= (1+\epsilon)^2 (1 + \epsilon \, |x|)^{\epsilon^{-1}}. \end{aligned}$$

□

Proof of Theorem 2.1-(c). Suppose $F \in \mathcal{D}(\Lambda)$ and $1-F$ is Γ varying so that

$$\lim_{n\to\infty} \frac{\overline{F}(b_n + x a_n)}{\overline{F}(b_n)} \quad = \quad e^{-x}, \;\; x \in \mathbb{R},$$

where the function $a_n = u(b_n) = \int_{b_n}^{r(F)} \overline{F}(s)ds/\overline{F}(b_n)$ is called an auxiliary function. From (1.1)

$$\lim_{n\to\infty} F^n(a_n x + b_n) \quad = \quad \Lambda(x), \; x \in \mathbb{R},$$

with $a_n = u(b_n)$ and $b_n = F^{\leftarrow}(1-\frac{1}{n}) = \inf\{x : F(x) > 1-\frac{1}{n}\}$, $n \geq 1$. From Theorem B.5, F satisfies the von Mises condition:

$$\lim_{n\to\infty} \frac{f(b_n)u(b_n)}{\overline{F}(b_n)} = 1. \tag{3.14}$$

Now, by Theorem B.6 implies the following density convergence on compact sets:

$$\lim_{n\to\infty} g_n(x) = \lambda(x), \; x \in K \subset \mathbb{R}, \tag{3.15}$$

where K is a compact set, and $g_n(x) = n\, a_n\, f(a_n x + b_n) F^{n-1}(a_n x + b_n)$.

We write

$$H_\beta(g_n) \quad = \frac{1}{(1-\beta)} \log\left[\int_{-v}^{v} (g_n(x))^\beta dx + \int_{-\infty}^{-v} (g_n(x))^\beta dx + \int_{v}^{\infty} (g_n(x))^\beta dx\right].$$

It is enough to show,

$$\lim_{v\to\infty}\lim_{n\to\infty}\left[\int_{-\infty}^{-v} (g_n(x))^\beta dx + \int_{v}^{\infty} (g_n(x))^\beta dx\right] = 0.$$

Set, $I_E(n,v) = \int_v^\infty (g_n(x))^{\beta-1} g_n(x) dx$, we have

$$0 < I_E(n,v) < J(n,\beta)\left(\frac{f(a_n v + b_n)}{f(b_n)}\right)^{\beta-1} (1 - F^n(a_n v + b_n)).$$

where, $J(n,\beta) = \left(n\bar{F}(b_n)\frac{f(b_n)a_n}{F(b_n)}\right)^{\beta-1} \to 1$ as $n \to \infty$ and, $0 \leq F(.) \leq 1$, for $\beta > 1$ and f is decreasing function then $\frac{f(a_n v + b_n)}{f(b_n)} < 1$ for $v \geq 1$. Therefore,

$$\lim_{v\to\infty}\lim_{n\to\infty} I_E(n,v) = 0. \tag{3.16}$$

Now, we choose ξ_n satisfying $-\log F(\xi_n) \simeq n^{-1/2}$. If $t_n = \frac{\xi_n - b_n}{a_n} \to c$, then $n^{1/2} \simeq -n \log F(a_n t_n + b_n) \to e^{-c}$, and this is contradict with $n^{1/2} \to \infty$. Therefore, $t_n = \frac{\xi_n - b_n}{a_n} \to -\infty$, as $\xi_n \to r(F)$ for large n.

We now decompose the integral,

$$I_F(n,v) = \int_{-\infty}^{t_n} (g_n(x))^\beta dx + \int_{t_n}^{-v} (g_n(x))^\beta dx = I_{F_1}(n) + I_{F_2}(n,v).$$

Set

$$\begin{aligned} 0 < I_{F_1}(n) \quad &< \quad n^\beta a_n^{\beta-1} F^{\beta(n-1)}(\xi_n) \int_{\mathbb{R}} (f(x))^\beta dx, \\ &\simeq \quad \frac{n^\beta a_n^{\beta-1}}{\exp\{n^{1/2}\beta\}} \exp\{\beta\, n^{-1/2}\} \int_{-\infty}^{\infty} (f(x))^\beta dx, \end{aligned}$$

where, $F^{n-1}(\xi_n) = \exp\Big\{(n-1)\log F(\xi_n)\Big\} \simeq \exp\{-n^{1/2} + n^{-1/2}\}$. From Theorem B.1, $a_n \in RV_0$ and from Lemma B.1, for $n > N$ given $\epsilon > 0$ and $\epsilon(n) < \epsilon$ then

$$a_n = c(n) \exp\Big\{\int_N^n \rho(t) t^{-1} dt\Big\} < (n/N)^\epsilon c,$$

Therefore,

$$0 < I_{F_1}(n) \quad \leq \quad \frac{cn^{\beta}(n/N)^{(\beta-1)\epsilon}}{\exp\{\beta(n^{1/2}-n^{-1/2})\}}\int_{-\infty}^{\infty}(f(s))^{\beta}\,ds.$$

If $\int_{-\infty}^{\infty}(f(x))^{\beta}dx < \infty$ then

$$\lim_{n\to\infty} I_{F_1}(n) = 0. \tag{3.17}$$

From (3.14), for given $\epsilon_1 > 0$ we have $f(a_nx+b_n) \leq -(1+\epsilon_1)\frac{\log F(a_nx+b_n)}{u(a_nx+b_n)}$ ultimately and from Theorem B.7 given $\epsilon_2 > 0$ and for $x < 0$ such that $a_nx+b_n \geq n_0$, $\frac{u(b_n)}{u(a_nx+b_n)} < \frac{1}{1-\epsilon_2}\left[\frac{-\log F(a_nx+b_n)}{-\log F(b_n)}\right]^{\epsilon_2}$ and for $\epsilon_3 > 0$ and for all $n \geq 1$ then $1-\epsilon_3 < \frac{n-1}{n}$, and from Lemma 3.3 for $\epsilon_4 > 0$ and $x < 0$ we have $n\bar{F}(a_nx+b_n) < (1+\epsilon_4)^2(1+\epsilon_4|x|)^{\epsilon_4^{-1}}$, and for $\epsilon_5 > 0$, $\frac{1}{F(b_n)} < \frac{n}{1-\epsilon_5}$. Therefore,

$$\begin{aligned}
g_n(x) &= na_nf(a_nx+b_n)F^{n-1}(a_nx+b_n),\\
&< -(1+\epsilon_1)\frac{u(b_n)}{u(a_nx+b_n)}n\log F(a_nx+b_n)\exp\Big\{\frac{n-1}{n}n\log F(a_nx+b_n)\Big\},\\
&< \frac{1+\epsilon_1}{(1-\epsilon_2)(1-\epsilon_5)^{1+\epsilon_2}}(-n\log F(a_nx+b_n))^{1+\epsilon_2}\exp\Big\{(1-\epsilon_3)n\log F(a_nx+b_n)\Big\},\\
&< \frac{(1+\epsilon_1)(1+\epsilon_4)^{2\frac{1+\epsilon_2}{\epsilon_4}}}{(1-\epsilon_2)(1-\epsilon_5)^{1+\epsilon_2}}(1+\epsilon_4|x|)^{\frac{1+\epsilon_2}{\epsilon_4}}\exp\Big\{-(1-\epsilon_3)(1+\epsilon_4)^2(1+\epsilon_4|x|)^{\epsilon_4^{-1}}\Big\},\\
&= c_1(1+\epsilon_4|x|)^{c_2}\exp\Big\{-c_3(1+\epsilon_4|x|)^{\epsilon_4^{-1}}\Big\},
\end{aligned}$$

where, $c_1 = \frac{(1+\epsilon_1)(1+\epsilon_4)^{2\frac{1+\epsilon_2}{\epsilon_4}}}{(1-\epsilon_2)(1-\epsilon_5)^{1+\epsilon_2}}$ and $c_2 = \frac{1+\epsilon_2}{\epsilon_4}$, and $c_3 = (1-\epsilon_3)(1+\epsilon_4)^2$. We define

$$h_1(x) = c_1(1+\epsilon_4|x|)^{c_2}\exp\Big\{-c_3(1+\epsilon_4|x|)^{\epsilon_4^{-1}}\Big\}. \tag{3.18}$$

And

$$\int_{-\infty}^{0}h_1(x)dx \quad = \quad \int_{-\infty}^{0}c_1(1+\epsilon_4|x|)^{c_2}\exp\Big\{-c_3(1+\epsilon_4|x|)^{\epsilon_4^{-1}}\Big\}dx.$$

Putting, $c_3(1+\epsilon_4|x|)^{\epsilon_4^{-1}} = y$, and $-c_3(1+\epsilon_4|x|)^{\epsilon_4^{-1}-1}dx = dy$ we have,

$$\int_{-\infty}^{0}h_1(x)dx \quad = \quad c_1\left(\frac{1}{c_3}\right)^{\epsilon_4(c_2+1)}\int_{c_3}^{\infty}y^{\epsilon_4(c_2+1)-1}e^{-y}dy < \infty.$$

Similarly, from (3.14) and from Theorem B.7 given $\epsilon_2 > 0$ and for $x > 0$ such that $a_nx+b_n \geq n_0$, $\frac{u(b_n)}{u(a_nx+b_n)} < \frac{1}{1-\epsilon_2}\left[\frac{-\log F(a_nx+b_n)}{-\log F(b_n)}\right]^{-\epsilon_2}$ and from Lemma 3.3 for $\epsilon_4 > 0$ and $x > 0$ we have $n\bar{F}(a_nx+b_n) < (1+\epsilon_4)^2(1-\epsilon_4x)^{\epsilon_4^{-1}}$, we have

$$\begin{aligned}
g_n(x) &= na_nf(a_nx+b_n)F^{n-1}(a_nx+b_n),\\
&< -(1+\epsilon_1)\frac{u(b_n)}{u(a_nx+b_n)}n\log F(a_nx+b_n)\exp\Big\{\frac{n-1}{n}n\log F(a_nx+b_n)\Big\},\\
&< \frac{1+\epsilon_1}{(1-\epsilon_2)(1-\epsilon_5)^{1-\epsilon_2}}(-n\log F(a_nx+b_n))^{1-\epsilon_2}\exp\Big\{(1-\epsilon_3)n\log F(a_nx+b_n)\Big\},\\
&< \frac{(1+\epsilon_1)(1+\epsilon_4)^{2\frac{1-\epsilon_2}{\epsilon_4}}}{(1-\epsilon_2)(1-\epsilon_5)^{1-\epsilon_2}}(1-\epsilon_4x)^{\frac{1-\epsilon_2}{\epsilon_4}}\exp\Big\{-(1-\epsilon_3)(1+\epsilon_4)^2(1-\epsilon_4x)^{\epsilon_4^{-1}}\Big\},\\
&= c_1(1-\epsilon_4x)^{c_2}\exp\Big\{-c_3(1-\epsilon_4x)^{\epsilon_4^{-1}}\Big\},
\end{aligned}$$

where, $c_1 = \frac{(1+\epsilon_1)(1+\epsilon_4)^{2\frac{1-\epsilon_2}{\epsilon_4}}}{(1-\epsilon_2)(1-\epsilon_5)^{1-\epsilon_2}}$ and $c_2 = \frac{1-\epsilon_2}{\epsilon_4}$, and $c_3 = (1-\epsilon_3)(1+\epsilon_4)^2$. We define

$$h_2(x) = c_1(1-\epsilon_4 x)^{c_2} \exp\Big\{ - c_3(1-\epsilon_4 x)^{\epsilon_4^{-1}}\Big\}. \tag{3.19}$$

And

$$\int_0^\infty h_2(x)dx \quad = \quad \int_0^\infty c_1(1-\epsilon_4 x)^{c_2} \exp\Big\{ - c_3(1-\epsilon_4 x)^{\epsilon_4^{-1}}\Big\}dx.$$

Putting $c_3(1-\epsilon_4 x)^{\epsilon_4^{-1}} = y$, and $-c_3(1-\epsilon_4 x)^{\epsilon_4^{-1}-1}dx = dy$ we have,

$$\int_0^\infty h_2(x)dx \quad = \quad c_1\left(\frac{1}{c_3}\right)^{\epsilon_4(c_2+1)} \int_0^{c_3} y^{\epsilon_4(c_2+1)-1}\, e^{-y}dy < \infty.$$

where, $\epsilon_4 > \epsilon_2$.
Now, we set $I_{F_2}(n,v) = \int_{t_n}^{-v} (g_n(x))^\beta\, dx$. From (3.18) for large n, we have $g_n(x) < h_1(x)$, then

$$I_{F_2}(n,v) \quad < \quad \int_{-\infty}^{-v} c_1^\beta (1+\epsilon_4\,|x|)^{\beta c_2} \exp\Big\{ - c_3\beta(1+\epsilon_4\,|x|)^{\epsilon_4^{-1}}\Big\}dx. \tag{3.20}$$

Putting $c_3\beta(1+\epsilon_4\,|x|)^{\epsilon_4^{-1}} = y$, and $-c_3\beta(1+\epsilon_4\,|x|)^{\epsilon_4^{-1}-1}dx = dy$ we have,

$$0 < I_{F_2}(n,v) \quad < \quad c_1^\beta \left(\frac{1}{c_3}\right)^{\epsilon_4(c_2\beta+1)} \int_{c_3\beta(1+\epsilon_4 v)^{\frac{1}{\epsilon_4}}}^\infty y^{\epsilon_4(c_2\beta+1)-1}\, e^{-y}dy.$$

Therefore,

$$\lim_{v\to\infty}\lim_{n\to\infty} I_{F_2}(n,v) = 0. \tag{3.21}$$

From, (3.17) and (3.21)

$$\lim_{v\to\infty}\lim_{n\to\infty} I_F(n,v) = 0. \tag{3.22}$$

Set $I_G(n,v) = \int_{-v}^{v} (g_n(x))^\beta\, dx$. From (3.18) and (3.19) for large n, we have $g_n(x) < h_1(x) + h_2(x)$, for and $\int_{-v}^0 h_1(x)dx + \int_0^v h_2(x)dx < \infty$, and using (3.15) $\lim_{n\to\infty} g_n(x) = \lambda(x)$, for $x \in [-v,v]$ by DCT,

$$\lim_{v\to\infty}\lim_{n\to\infty} \int_{-v}^{v} (g_n(x))^\beta\, dx \quad = \quad \int_{-\infty}^{\infty} (\lambda(x))^\beta\, dx. \tag{3.23}$$

From, (3.16), (3.22) and (3.23),

$$\lim_{n\to\infty} H_\beta(g_n) = H_\beta(\lambda).$$

□

Appendix A.

Lemma A.1. The Rényi entropy of

(i) Fréchet law:

$$H_\beta(\phi_\alpha) = -\log\alpha + \frac{\alpha+1}{\alpha}\log\beta - \frac{1}{1-\beta}\left(\log\beta - \log\Gamma\left(\frac{\alpha+1}{\alpha}(\beta-1)+1\right)\right);$$

where, $\frac{1}{\alpha+1} < \beta$.

(ii) Weibull law:

$$H_\beta(\psi_\alpha) = -\log\alpha + \frac{\alpha-1}{\alpha}\log\beta - \frac{1}{1-\beta}\left(\log\beta - \log\Gamma\left(\frac{\alpha-1}{\alpha}(\beta-1)+1\right)\right);$$

where, $\max\left(0, \frac{\beta-1}{\beta}\right) < \alpha$, for $\beta > 0$.

(iii) Gumbel law:

$$H_\beta(\lambda) = \frac{1}{1-\beta}\log\frac{\Gamma(\beta)}{\beta^\beta};$$

where, $\beta > 0$.

Proof. (i) The Rényi entropy of Fréchet distribution is

$$H_\beta(\phi_\alpha) = \frac{1}{1-\beta}\log\int_0^\infty \left(\alpha x^{-\alpha-1}e^{-x^{-\alpha}}\right)^\beta dx.$$

Putting, $\beta x^{-\alpha} = u$, $-\beta\alpha x^{-\alpha-1}dx = du$,

$$\begin{aligned} H_\beta(\phi_\alpha) &= \frac{1}{1-\beta}\log\int_0^\infty \alpha^{\beta-1}u^{(\beta-1)(\frac{\alpha+1}{\alpha})}\beta^{-((\beta-1)(\frac{\alpha+1}{\alpha})+1)}e^{-u}du, \\ &= \frac{1}{1-\beta}\left((\beta-1)\log\alpha - \left((\beta-1)\frac{\alpha+1}{\alpha}+1\right)\log\beta + \log\Gamma\left(\frac{\alpha+1}{\alpha}(\beta-1)+1\right)\right). \end{aligned}$$

where, $\frac{\alpha+1}{\alpha}(\beta-1)+1 > 0$, so $\frac{1}{\beta} < \alpha+1$.
(ii) The Rényi entropy of Weibull distribution

$$H_\beta(\psi_\alpha) = \frac{1}{1-\beta}\log\int_{-\infty}^0 \left(\alpha(-x)^{\alpha-1}e^{-(-x)^\alpha}\right)^\beta dx.$$

Putting, $\beta(-x)^\alpha = u$, $-\beta\alpha(-x)^{\alpha-1}dx = du$,

$$\begin{aligned} H_\beta(\psi_\alpha) &= \frac{1}{1-\beta}\log\int_0^\infty \alpha^{\beta-1}u^{(\beta-1)(\frac{\alpha-1}{\alpha})}\beta^{-((\beta-1)(\frac{\alpha-1}{\alpha})+1)}e^{-u}du, \\ &= \frac{1}{1-\beta}\left((\beta-1)\log\alpha - \left((\beta-1)\frac{\alpha-1}{\alpha}+1\right)\log\beta + \log\Gamma\left(\frac{\alpha-1}{\alpha}(\beta-1)+1\right)\right), \end{aligned}$$

where, $\frac{\alpha-1}{\alpha}(\beta-1)+1 > 0$. If $\beta > 1$ then $\alpha > \frac{\beta-1}{\beta}$, and for $\beta < 1$, we have $0 < \alpha$, therefore, $\max\left(0, \frac{\beta-1}{\beta}\right) < \alpha$, for all $\beta > 0$.
(iii) The Rényi entropy of Gumbel distribution

$$H_\beta(\lambda) = \frac{1}{1-\beta}\log\int_{-\infty}^\infty \left(e^{-x}e^{-e^{-x}}\right)^\beta dx,$$

Taking $\frac{u}{\beta} = e^{-x}$ and $\frac{du}{\beta} = -e^{-x}dx$

$$\begin{aligned} H_\beta(\lambda) &= \frac{1}{1-\beta}\log\int_0^\infty u^{\beta-1}e^{-u}\beta^{-\beta}du, \\ &= \frac{1}{1-\beta}\left(\log\Gamma(\beta) - \beta\log\beta\right). \end{aligned}$$

where, $\beta > 0$. □

Lemma A.2. If $Y = \frac{X-b}{a}$, for $b \in \mathbb{R}$ and $a > 0$, then the Rényi's entropy of Y is given by

$$H_\beta(f_Y) = -\log a + H_\beta(f_X).$$

Proof. We have $F_Y(y) = \Pr(X \leq ay+b) = F_X(ay+b)$, and $f_Y(y) = af_X(ay+b)$, so that from (1.2),

$$\begin{aligned} H_\beta(f_Y) &= \frac{1}{1-\beta}\log\int_{-\infty}^{\infty}(af_X(ay+b))^\beta\,dy = \frac{1}{1-\beta}\log\int_{-\infty}^{\infty}f_X^\beta(z)a^{\beta-1}dz, \\ &= -\log a + H_\beta(f_X). \end{aligned}$$

□

Appendix B.

Definition 1. (Definition, Page 27, Resnick (1987)) A nonegative, nondecreasing function $V(x)$ defined on a semi infinite interval (z,∞) is Π varying (written $V \in \Pi$) if there exist functions $a(t) > 0$, $b(t) \in \mathbb{R}$ such that for $x > 0$

$$\lim_{t\to\infty}\frac{V(tx)-b(t)}{a(t)} = \log x.$$

Theorem B.1. (Proposition 0.12, Resnick (1987)) If $V \in \Pi$ with auxiliary function $a(t)$ then $a(\cdot) \in RV_0$.

Theorem B.2. (Proposition 0.8, Resnick (1987)) Suppose $U \in RV_\rho$, $\rho \in \mathbb{R}$. Take $\epsilon > 0$. Then there exists t_0 such that for $x \geq 1$ and $t \geq t_0$

$$(1-\epsilon)x^{\rho-\epsilon} < \frac{U(tx)}{U(t)} < (1+\epsilon)x^{\rho+\epsilon}.$$

Lemma B.1. (Corollary, Page 17, Resnick (1987)) If L is slowly varying iff L can be represented as

$$L(x) = c(x)\exp\Big\{\int_1^x t^{-1}\epsilon(t)dt\Big\}, \tag{B.1}$$

for $x > 0$ and $\lim_{x\to\infty} c(x) = c$ and $\lim_{t\to\infty}\epsilon(t) = 0$.

Remark B.1. (Remark, Page 19, Resnick (1987)) If $U \in RV_\rho$ then U has representation

$$U(x) = c(x)\exp\Big\{\int_1^x t^{-1}\rho(t)dt\Big\}, \tag{B.2}$$

where, $\lim_{x\to\infty} c(x) = c$ and $\lim_{t\to\infty}\rho(t) = \rho$.

Theorem B.3. (Proposition 2.1, Resnick (1987)) Let $F \in \mathcal{D}(G)$.

(i) If $G = \Phi_\alpha$, then $a_n = (1/(1-F))^{\leftarrow}(n)$, $b_n = 0$, and if for some integer $0 < k < \alpha$,

$$\int_{-\infty}^{0} |\,x\,|^k\,F(dx) < \infty,$$

then $\lim_{n\to\infty} E\left(\frac{M_n}{a_n}\right)^k = \int_0^\infty x^k\Phi_\alpha(dx) = \Gamma\left(1-\frac{k}{\alpha}\right)$.

(ii) If $G = \Psi_\alpha$, then $\delta_n = r(F) - (1/(1-F))^{\leftarrow}(n)$, $r(F) < \infty$, and if for some integer $k > 0$,

$$\int_{-\infty}^{r(F)} |\,x\,|^k\,F(dx) < \infty,$$

then $\lim_{n\to\infty} E\left(\frac{M_n - r(F)}{\delta_n}\right)^k = \int_{-\infty}^0 x^k\Psi_\alpha(dx) = (-1)^k\Gamma\left(1+\frac{k}{\alpha}\right)$.

(iii) If $G = \Lambda$, then $b_n = (1/(1-F))^{\leftarrow}(n)$, $a_n = u(b_n)$, and if for some integer $k > 0$,

$$\int_{-\infty}^{0} |\ x\ |^k\ F(dx) < \infty,$$

then $\lim_{n\to\infty} E\left(\frac{M_n - b_n}{a_n}\right)^k = \int_{-\infty}^{\infty} x^k \Lambda(dx) = (-1)^k \Gamma^{(k)}(1)$, where $\Gamma^{(k)}(1)$ is the k-th derivative of the Gamma function at $x = 1$.

Theorem B.4. (Proposition 1.15 and 1.16, Resnick (1987))
(i) Suppose that distribution function F is absolutely continuous with density f which is eventually positive.

(a) If for some $\alpha > 0$

$$\lim_{x\to\infty} \frac{xf(x)}{\overline{F}(x)} = \alpha, \tag{B.3}$$

then $F \in \mathcal{D}(\Phi_\alpha)$.

(b) If f is nonincreasing and $F \in \mathcal{D}(\Phi_\alpha)$ then (B.3) holds.
(ii) Suppose F has right endpoint $r(F)$ finite and density f positive in a left neighbourhood of $r(F)$.

(a) If for some $\alpha > 0$

$$\lim_{x\to r(F)} \frac{(r(F) - x)f(x)}{\overline{F}(x)} = \alpha \tag{B.4}$$

then $F \in \mathcal{D}(\Psi_\alpha)$.

(b) If f is nonincreasing and $F \in \mathcal{D}(\Psi_\alpha)$ then (B.4) holds.

Theorem B.5. (Proposition 1.17, Resnick (1987)) Let F be absolutely continuous in a left neighborhood of $r(F)$ with density f. If

$$\lim_{x\uparrow r(F)} f(x) \int_x^{r(F)} \overline{F}(t)dt/\overline{F}(x)^2 = 1, \tag{B.5}$$

then $F \in \mathcal{D}(\Lambda)$. In this case we may take,

$$u(x) = \int_x^{r(F)} \overline{F}(t)dt/\overline{F}(x),\ b_n = F^{\leftarrow}(1 - 1/n),\ \ a_n = u(b_n).$$

Theorem B.6. (Theorem 2.5, Resnick (1987)), Suppose that F is absolutely continuous with pdf f. If $F \in \mathcal{D}(G)$ and

(i) $G = \Phi_\alpha$, then $g_n(x) \to \phi_\alpha(x)$ locally uniformly on $(0, \infty)$ iff (B.3) holds;
(ii) $G = \Psi_\alpha$, then $g_n(x) \to \psi_\alpha(x)$ locally uniformly on $(-\infty, 0)$ iff (B.4) holds;
(iii) $G = \Lambda$, then $g_n(x) \to \lambda(x)$ locally uniformly on $\mathbb{R}$ iff (B.5) holds.

Theorem B.7. (Lemma 2, De Haan and Resnick (1982)) Suppose $F \in \mathcal{D}(\Lambda)$ with auxiliary function u and $\epsilon > 0$. There exists a t_0 such that for $x \geq 0$, $t \geq t_0$

$$(1-\epsilon)\left[\frac{-\log F(t)}{-\log F(t + xu(t))}\right]^{-\epsilon} \leq \frac{u(t + xu(t))}{u(t)} \leq (1+\epsilon)\left[\frac{-\log F(t)}{-\log F(t + xu(t))}\right]^{\epsilon},$$

and for $x < 0$, $t + xu(t) \geq t_0$

$$(1-\epsilon)\left[\frac{-\log F(t + xu(t))}{-\log F(t)}\right]^{-\epsilon} \leq \frac{u(t + xu(t))}{u(t)} \leq (1+\epsilon)\left[\frac{-\log F(t + xu(t))}{-\log F(t)}\right]^{\epsilon}.$$

Theorem B.8. (Corollary, Balkema and De Haan (1972)) A distribution function $F \in \mathcal{D}(\Lambda)$ if and only if there exist a positive function c satisfying $\lim_{x \to r(F)} c(x) = 1$ and a positive differentiable function $u(t)$ satisfying $\lim_{x \to r(F)} u'(x) = 0$ such that

$$\bar{F}(x) = c(x) \exp\Big\{ - \int_{-\infty}^{x} \frac{dt}{u(t)} \Big\} \text{ for } x < r(F).$$

References

[1] Artstein, S. and Ball, K.M. and Barthe, F. and Naor, A., (2004), *On the rate of convergence in the entropic central limit theorem*, *Probability Theory Related Fields*, Vol. 129, Pages 381-390.

[2] Barron A.R., (1986), *Entropy and the Central Limit Theorem, The Annals of Probability*, Vol. 14, No. 1, Pages 336-342.

[3] Balkema, A. A. and De Haan, L., (1972), *On R. von Mises condition for the domain of attraction of* $\exp(-e^{-x})$*, The Annals of Mathematical Statistics*, Vol. 43, No. 4, Pages 1352-1354.

[4] Brown, L. D., (1982), *A proof of the central limit theorem motivated by the Cramer-Rao inequality*, *Statistics and probability: Essays in Honour of C.R. Rao*, Pages 141-148.

[5] Cui, Hongfei and Ding, Yiming, (2010), *The convergence of the Rényi entropy of the normalized sums of IID random variables, Statistics and Probability Letters*, Vol. 80, Pages 1167-1173.

[6] De Haan, Laurens and Resnick, Sidney I., (1982), *Local limit theorems for sample extremes, The Annals of Probability*, Vol. 10, No. 2, Pages 396-413.

[7] Embrechts P. and Klüppelberg C. and Mikosch, T., (1997), *Modelling Extremal Events for Insurance and Finance,* Springer Verlag.

[8] Galambos J., (1987), *The Asymptotic Theory of Extreme Order Statistics,* Krieger Pub. Co.

[9] Johnson O., (2006), *Information Theory and the Central Limit Theorem,* Imperical College Press.

[10] Johnson, O. and Barron, A., (2004), *Fisher information inequalities and the central limit theorem, Probab. Theory Relat. Fields*, Vol. 129, Pages 391-409.

[11] Linnik, J. V., (1959), *An information theoretic proof of the central limit theorem with Lindeberg conditions*, *Theory Probability Application*, Vol. 4, Pages 288-299.

[12] Resnick Sidney I., (1987), *Extreme Values, Regular Variation, and Point Processes,* Springer Verlag.

[13] Renyi, A., (1961), *On Measures of Entropy and Information, Proceedings of the Fourth Berkeley Symposium on Mathematics, Statistics and Probability, University of California Press*, Vol. 1, Pages 547-561.

[14] Saeb, A., (2014), *Rates of convergence for Rényi entropy in extreme value theory,* (Preprint).

[15] Shimizu, R., (1975), *On fisher's amount of information for location family*, *Statistical distributions in scientific work*, Vol. 3, Pages 305-312.

[16] Takano, Seiji, (1987), *Convergence entropy in the central limit theorem*, *Yokohama Mathematical Journal*, Vol. 35.